

## Stress MIBI Scan medication instructions:

<u>Generic Name</u>	<u>Brand Name</u>
<b>Type 1</b> Stop for 72 hours (3 full days) before day of Exercise Study	
Dipyridamole	Asasantin, Persantin
Sildenafil	Viagra
<b>TYPE 2</b> Stop for 48 hours (2 full days) before day of Exercise Study	
<b>Do not stop Sotolol. Carvedilol or Bisoprolol</b>	
Atenolol	Tenormin, Noten, Tensig, Atehexal
Metoprolol	Betaloc, Metohexal, Minax
Oxprenolol	Corbeton, Trasicor
Pindolol	Barbloc, Visken
Propranolol	Inderal, Deralin, Cardinol
<b>TYPE 3</b> Stop for 24 hours (1 full day) before day of Exercise Study	
Verapamil	Isoptin, Veracaps, Ampec, Cordilox, Tarka
Nicorandil	Ikorel
Diltiazem	Vasocardol, Cardizem, Dilzem, Coras, Diltahexal
Felodipine	Felodur, Plendil, Agon, Triasyn, Felodil
Isosorbide	Isordil, Duride, Imdur, Monodur
Glyceryl Trinitrate	Nitro-bid, Nitro-dur, Transiderm Nitradisc
Lercanidipine	Zanidip, Zan-extra
<b>OTHER</b> Stop on day of Exercise Study	
Amlodipine	Norvasc, Caduet, Perivasc, Amlo, Ozlopid, Exforge, Amiotrust
Nicotine Patches	
Nifedipine	Adalat, Adalat Orus, Addos, Adefin